

DECREASING DISPARITIES IN DEPRESSION CARE – A Core Curriculum for Change

REFERENCES

Aina Y, Susman JL. Understanding comorbidity with depression and anxiety disorders. *J Am Osteopath Assoc*. 2006;106(5 Suppl 2):9-14.

American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, 4th Edition, Text Revision (DSM-IV-TR)*. Washington, DC: American Psychiatric Association; 2000.

American Psychiatric Association. Practice Guideline for the Treatment of Patients with Bipolar Disorders. 2002. Available at: http://www.psych.org/psych_pract/treatg/pg/prac_guide.cfm. Accessed September 6, 2006.

Arias E, Anderson RN, Hsiang-Ching K, Murphy SL, Kochanek KD. *Deaths: Final Data for 2001. National Vital Statistics Reports*. Hyattsville, Maryland: National Center for Health Statistics. 2003;52(3). Available at: http://www.cdc.gov/nchs/data/nvsr/nvsr52/nvsr52_03.pdf. Accessed October 2, 2006.

Boulware LE, Cooper LA, Ratner LE, LaVeist TA, Powe NR. Race and trust in the health care system. *Public Health Rep*. 2003;118:358-365.

Centers for Disease Control and Prevention. *Morbidity and Mortality Weekly Report*. Self-reported frequent mental distress among adults – United States, 1993-1996. 1998;47(16):325-331. Atlanta, GA: CDC. Available at: <http://www.cdc.gov/mmwr/preview/mmwrhtml/00052469.htm>. Accessed October 25, 2006.

Chapman DP, Perry GS, Strine TW. The vital link between chronic disease and depressive disorders. *Prev Chronic Dis*. 2005;2(1):A14.

Ferketich AK, Schwartzbaum JA, Frid DJ, Moeschberger ML. Depression as an antecedent to heart disease among women and men in the NHANES I study. National Health and Nutrition Examination Survey. *Arch Intern Med*. 2000;160(9):1261-8.

Frasure-Smith N, Lesperance F, Talajic M. Depression following myocardial infarction: impact on 6-month survival. *JAMA*. 1993;270:1819-25.

Goldberg, RJ. *Practical Guide to the Care of the Psychiatric Patient*. 2nd ed. St. Louis, MO: Mosby; 1998;229.

Grant BF, Stinson FS, Dawson DA, et al. Prevalence and co-occurrence of substance use disorders and independent mood and anxiety disorders. *Arch Gen Psych*. 2004;61:807-816.

Greenberg PE, Kessler RC, Birnbaum HG, Leong SA, Lowe SW, Berglund PA, Corey-Lisle PK. The economic burden of depression in the United States: How did it change between 1990 and 2000? *J Clin Psychiatry*. 2003;64(12):1465-75.

Katon W, Schulberg H. Epidemiology of depression in primary care. *Gen Hosp Psychiatry*. 1992;14:237-247.

Kessler RC, Wai Tat Chiu AM, Demler MA, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*. 2005;62:617-709.

Kroenke K. The interface between physical and psychological symptoms. *Primary Care Companion J Clin Psychiatry*. 2003;5(suppl 7):11-18.

Landheim AS, Bakken K, Vaglum P. Impact of comorbid psychiatric disorders on the outcome of substance abusers: a six year prospective follow-up in two Norwegian counties. *BMC Psychiatry*. 2006;6:44-54.

Lustman PJ, Anderson RJ, Freedland KE, de Groot M, Carney RM, Clouse RE. Depression and poor glycemetic control: a meta-analytic review of the literature. *Diabetes Care*. 2000;23(7):934-42.

National Center for Health Statistics. *Health, United States, 2005*. Available at: [http://www.cdc.gov/nchs/data/05.pdf#046](http://www.cdc.gov/nchs/data/hus/05.pdf#046). Accessed December 11, 2006.

National Institute of Mental Health. *In Harm's Way: Suicide in America*. Bethesda (MD): National Institute of Mental Health, National Institutes of Health, US Dept of Health and Human Services. Printed January 2001/ Revised May 2003. (NIH Publication No. 03-4594). Available at: <http://www.nimh.nih.gov/publicat/harmsway.cfm>. Accessed September 1, 2006.

Regier DA, Narrow WE, Rae DS, Manderscheid RW, Locke BZ, Goodwin FK. The de facto US mental and addictive disorders service system: epidemiologic catchment area prospective 1-year prevalence rates of disorders and services. *Arch Gen Psychiatry*. 1993;50:85-94.

Rost K, Pyne JM, Dickinson LM, LoSasso AT. Cost-effectiveness of enhancing primary care depression management on an ongoing basis. *Ann Fam Med*. 2005; 3:7-14.

Saluja G, Iachan R, Scheidt PC, Overpeck MD, Sun W, Giedd JN. Prevalence of and risk factors for depressive symptoms among young adolescents. *Arch Pediatr Adolesc Med*. 2004;158:760-765.

Simon G, Ormel J, VonKorff M, Barlow W. Health care costs associated with depressive and anxiety disorders in primary care. *Am J Psychiatry*. 1995;152:352-7.

US Census Bureau. *Profiles of General Demographic Characteristics: 2000*. Washington, DC, US Census Bureau, May 2001. Available at: <http://www.census.gov/Press-Release/www/2001/demoprofile.html>. Accessed September 12, 2006.

US Department of Health and Human Services. *Mental Health: A Report of the Surgeon General*. Rockville, MD: US Dept. of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services; 1999.

US Department of Health and Human Services. *Mental Health: Culture, Race, and Ethnicity – A Supplement to Mental Health: A Report of the Surgeon General*. Rockville, MD: US Dept. of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services; 2001.

US Department of Health and Human Services. *Mental Health: Culture, Race, and Ethnicity: Fact Sheets*. 2001. Available at: <http://mentalhealth.samhsa.gov/cre/>. Accessed December 11, 2006.

Üstün TB, Ayuso-Mateos JL, Chatterji S, Mathers C, Murray CJL. Global burden of depressive disorders in the year 2000. *Br J Psychiatry*. (2004) 184: 386-392.